

# Alliance Seeds

"Seeding the future together"

## TOMATO – MAVIRI F1

### INDETERMINATE SALADETTE

#### DESCRIPTION:

#### PLANT

- **Vigour** - Vigorous Plant with short internodes
- **Habit** - Indeterminate
- **Leaf** - Good foliage cover

#### FRUIT

- **Shape** - Elongated saladette with thick wall
- **Colour** - Uniform ripening, shiny red colour
- **Size** - 120–140g
- **Firmness** - good



#### RESISTANCES:

- Resistant (HR) Vd:1 (*Verticillium dahlia*, race 1) / Va:1 (*Verticillium albo-atrum*, race, 1) / Fol 1,2,3 (Fusarium Wilt races 1,2,3) / ToMV (Tomato Mosaic Virus) / Lt (Powdery Mildew)
- Resistant (IR) M (Nematodes) *Ma, Mi, Mj* Root Knot / TSWV (Tomato Spotted Wilt Virus)

#### PRODUCTION SLOT:

- Mainly protected crops

#### RECOMMENDATIONS:

- 12 500-15 000 double stem/ha open field
- 30 000 stems/ha protected cropping

#### ADVANTAGES:

- Vigorous plant with short internodes
- Concentrated setting, keeping size at top.
- Uniform fruit quality (shiny colour, deep red)
- Large fruit size with good fruit firmness
- Resistance to F3, Lt and TSWV

TEL: (013) 750 0575  
FAX: (013) 750 0656  
info@allianceseeds.co.za



#### IMPORTANT

The descriptions, advice, suggestions and vegetation cycles that may be presented herein are aimed at experienced professionals and are offered in all good faith, for purely informational purposes, and shall not therefore, under any circumstances, be held to be exhaustive, be taken as any form of guarantee of harvest, prejudice specific factors or circumstances (either current or future), and more generally, form of any kind of contractual undertaking whatsoever.

The user must first and foremost ensure that his local geographical conditions, his planned growing period, his soil, the means at his disposal (such as technical knowledge and experience and cultural techniques and operations) and his equipment, and more generally his agronomical, climatic, sanitary, environmental and economic context are suitable for the crops, techniques and varieties that are presented herein.

The varieties illustrated in this publication were photographed in favourable conditions and no guarantee can be provided that results will be identical under different conditions